

PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	for						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

Display	Abstract	Show: 20	Sort	Send to	Text
---------	----------	----------	------	---------	------

Entrez PubMed

1: Hum Immunol 1999 Jul;60(7):583-90

[Full-Text Articles](#)

NOD background genes influence T cell responses to GAD 65 in HLA-DQ8 transgenic mice.

Abraham RS, Wilson SB, de Souza NF Jr, Strominger JL, Munn SR, David CS.

PubMed Services

Department of Immunology, Division of Transplantation Surgery, Mayo Clinic, Rochester, MN 55905, USA.

Related Resources

The major histocompatibility complex (MHC) genes play a significant role in the predisposition to insulin-dependent diabetes mellitus or type 1 diabetes. HLA-DQ8 (DQB1*0302, DQA 1*0301) genes have been shown to have the highest relative risk for human type 1 diabetes. To develop a "humanized" mouse model of diabetes, HLA-DQ8 was transgenically expressed in mice lacking endogenous class II genes. Since non-MHC background genes of the NOD influence the disease process, AP^u/DQ8 mice were mated with the NOD strain and backcrossed to generate Abeta degree/DQ8/NOD mice. These mice have DQ8 as the sole MHC class II restriction element with NOD background genes at the N 2 generation. The DQ8 transgenic mice were used to identify T cell epitopes on glutamic acid decarboxylase (GAD 65), an important putative autoantigen in type 1 diabetes. The NOD background genes strongly influenced antigen processing, that is, different T cell epitopes were generated from the processing of GAD 65 in vivo in the Abeta degree/DQ8 and in the Abeta degree/DQ8/NOD mice.

PMID: 10426275 [PubMed - indexed for MEDLINE]


Display	Abstract	Show: 20	Sort	Send to	Text
---------	----------	----------	------	---------	------

[Write to the Help Desk](#)
[NCBI | NLM | NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act | Disclaimer](#)

Mar 6 2003 10:50:17

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI,TDBD; THES=ASSIGNEE; PLUR=YES; OP=AND</i>			
<u>L14</u>	Abraham-roshini-S\$.in.	0	<u>L14</u>
<u>L13</u>	David-chella-S\$.in.	1	<u>L13</u>
<u>L12</u>	(transgenic adj mouse) same (DR3)	6	<u>L12</u>
<u>L11</u>	(transgenic adj mouse) same (I-Ealpha)	1	<u>L11</u>
<u>L10</u>	L9 and (DRab and DQab)	2	<u>L10</u>
<u>L9</u>	L8 and (DR3)	27	<u>L9</u>
<u>L8</u>	L2 or L5 or L6 or L7	1637	<u>L8</u>
<u>L7</u>	(SCID adj (mouse or mice))	1630	<u>L7</u>
<u>L6</u>	((CD3 or Ig) adj knockout)	2	<u>L6</u>
<u>L5</u>	L4 and (TCR adj knockout)	1	<u>L5</u>
<u>L4</u>	(transgenic adj mouse) same (TCR)	129	<u>L4</u>
<u>L3</u>	(transgenic adj mouse) same (RAG-2)	12	<u>L3</u>
<u>L2</u>	(transgenic adj mouse) same (RAG-?)	13	<u>L2</u>
<u>L1</u>	Huang-manley.in.	1	<u>L1</u>

END OF SEARCH HISTORY



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	▼ for						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

☐ 1: Hum Immunol 2002 Nov;63(11):987-99

[Related Articles, Links](#)

Entrez PubMed

[Full-Text Article](#)

Modulation of insulinitis and type 1 diabetes by transgenic HLA-DR3 and DQ8 in NOD mice lacking endogenous MHC class II.

PubMed
Services

Kudva YC, Rajagopalan G, Raju R, Abraham RS, Smart M, Hanson J, David CS.

Division of Endocrinology and Metabolism, Mayo Clinic, Rochester, MN, USA.

Related
Resources

To evaluate the contributions of DR3 and DQ8 to the etiopathogenesis of type 1 diabetes in a diabetes-predisposing milieu, we developed human leukocyte antigen (HLA) transgenic mice on the nonobese diabetic (NOD) background in the absence of the endogenous class II molecule, I-A(g7) and studied the incidence of both spontaneous and experimental (induced) autoimmune diabetes. Transgenic expression of HLA-DR3 and -DQ8 (either alone or in combination) did not confer susceptibility to spontaneous or cyclophosphamide-induced type 1 diabetes. Expression of I-A(g7) was mandatory for development of spontaneous or cyclophosphamide-induced diabetes. However, multiple low doses of streptozotocin could induce diabetes in all groups of mice independent of the class II molecules expressed. In unmanipulated mice, only islets from I-A(g7+/+) mice revealed significant intra-islet infiltration. Although a characteristic peri-insulitis/peri-ductulitis was present in Abeta(0)/NOD mice, islets from DR3, DQ8 and DR3 x DQ8 double transgenic mice demonstrated significantly less infiltration. In conclusion, transgenic expression of HLA-DR3 and -DQ8 associated with predisposition to type 1 diabetes alone is not sufficient to induce spontaneous diabetes in NOD mice lacking endogenous class II molecules.

PMID: 12392851 [PubMed - in process]

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)